كلية الحاسبات والذكاء الإصطناعي

BS111

Probability and Statistics

Lecture 03

Dr. Ahmed Hagag

Faculty of Computers and Artificial Intelligence
Benha University

Spring 2019-2020

Chapter 1: Probability

- Sample Space.
- Events.
- Counting Techniques.
- Probability of an Event.
- Additive Rules.
- Conditional Probability.
- Independence, and the Product Rule.
- Bayes' Rule.

Probability of an Event

Axioms of Probability:

S Is a sample space, A is an event
$A \subseteq S$

$$\boldsymbol{P(S) = 1}$$

$$\boldsymbol{P(\emptyset) = 0}$$

$$\boldsymbol{0 \leq P(A) \leq 1}$$

$$\boldsymbol{P(A') = 1 - P(A)}$$

Additive Rules (1/13)

If A and B are two events, then

$$P(A \cup B) = P(A) + P(B) - P(A \cap B).$$



Additive Rules (2/13)

For three events A, B, and C,

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C).$$

Additive Rules (3/13)

If A and B are mutually exclusive, then

$$P(A \cup B) = P(A) + P(B).$$



Additive Rules (4/13)

If $A_1, A_2, \ldots, A_n$ are mutually exclusive, then

$$P(A_1 \cup A_2 \cup \cdots \cup A_n) = P(A_1) + P(A_2) + \cdots + P(A_n).$$

Additive Rules (5/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

1) $P(A')$
2) $P(A \cup B)$
3) $P(A' \cap B)$
4) $P[(A \cup B)']$

Additive Rules (6/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

1) $P(A') = 1 - P(A) = 1 - 0.3 = 0.7$

2) $P(A \cup B) = 0.3 + 0.2 - 0.1 = 0.4$

Additive Rules (7/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

3) $P(A' \cap B)$



Additive Rules (7/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

3) $P(A' \cap B)$



Additive Rules (7/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

3) $P(A' \cap B)$



Additive Rules (7/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

3) $P(A' \cap B)$

$= P(B) - P(A \cap B) = 0.2 - 0.1$

$= 0.1$



Additive Rules (8/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

4) $P[(A \cup B)']$

Additive Rules (8/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

4) $P[(A \cup B)'] = P(A' \cap B')$



Additive Rules (8/13)

Example4:

If $P(A) = 0.3$, $P(B) = 0.2$, $P(A \cap B) = 0.1$ determine the following probabilities:

4) $P[(A \cup B)'] = P(A' \cap B')$

$= 1 - P(A \cup B) = 1 - 0.4 = 0.6$



Additive Rules (9/13)

Example5:

What is the probability of getting a total of 7 or 11 when a pair of fair dice is tossed?

Additive Rules (10/13)

Example5:

What is the probability of getting a total of 7 or 11 when a pair of fair dice is tossed?

Solution:

Let A be the event that 7 occurs and B the event that 11 comes up.

$$S = \{11, 12, 13, 14, 15, 16, 21, 22, 23, 24, 25, 26,$$
$$31, 32, 33, 34, 35, 36, 41, 42, 43, 44, 45, 46, 51,$$
$$52, 53, 54, 55, 56, 61, 62, 63, 64, 65, 66\}$$

Additive Rules (11/13)

Example5:

What is the probability of getting a total of 7 or 11 when a pair of fair dice is tossed?

Solution:

Let A be the event that 7 occurs and B the event that 11 comes up.

$$P(A) = \frac{6}{36}$$

$S = \{11, 12, 13, 14, 15, 16, 21, 22, 23, 24, 25, 26,$
$31, 32, 33, 34, 35, 36, 41, 42, 43, 44, 45, 46, 51,$
$52, 53, 54, 55, 56, 61, 62, 63, 64, 65, 66\}$

Additive Rules (12/13)

Example5:

What is the probability of getting a total of 7 or 11 when a pair of fair dice is tossed?

Solution:

Let A be the event that 7 occurs and B the event that 11 comes up.

$$P(B) = \frac{2}{36}$$

$S = \{11, 12, 13, 14, 15, 16, 21, 22, 23, 24, 25, 26,$
$31, 32, 33, 34, 35, 36, 41, 42, 43, 44, 45, 46, 51,$
$52, 53, 54, 55, 56, 61, 62, 63, 64, 65, 66\}$

Additive Rules (13/13)

Example5:

What is the probability of getting a total of 7 or 11 when a pair of fair dice is tossed?

Solution:

The probability of getting a total of 7 or 11 $= P(A \cup B)$

The events A and B are mutually exclusive, since a total of 7 and 11 cannot both occur on the same toss. Therefore,

$$P(A \cup B) = P(A) + P(B) = \frac{6}{36} + \frac{2}{36} = \frac{8}{36} = \frac{2}{9}$$

Conditional Probability (1/31)

Introduction (1/2):

Sometimes probabilities need to be reevaluated as additional information becomes available. The probability of an event B under the knowledge that the outcome will be in event A is denoted as

$$P(B \mid A)$$

and this is called the conditional probability of B given A.

Conditional Probability (1/31)

Introduction (2/2):

The conditional probability of B, given A, denoted by $P(B \mid A)$, is defined by

$$P(B|A) = \frac{P(A \cap B)}{P(A)} \quad , \qquad \text{provided} \quad P(A) > 0$$

Conditional Probability (1/31)

Introduction (2/2):

The conditional probability of B, given A, denoted by $P(B \mid A)$, is defined by

$$P(B|A) = \frac{P(A \cap B)}{P(A)} \quad , \qquad \text{provided} \quad P(A) > 0$$

Equally Likely Outcomes

$$\frac{P(A \cap B)}{P(A)} = \frac{\text{number of outcomes in } A \cap B}{\text{number of outcomes in } A}$$

Conditional Probability (2/31)

Example1:

As an additional illustration, suppose that our sample space S is the population of adults in a small town who have completed the requirements for a college degree.
We shall categorize them according to gender and employment status. The data are given in the following.

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (3/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (3/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

$$P(M|E)$$

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (3/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

$$P(M|E) = \frac{P(M \cap E)}{P(E)}$$

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (4/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

$$P(M \cap E) = \frac{460}{900}$$

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (5/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

$$P(E) = \frac{600}{900}$$

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (5/31)

Example1:

One of these individuals is to be selected at random. What is the probability that he is a male and given that he is employed?

$$P(M|E) = \frac{460/900}{600/900} = \frac{460}{600}$$

Categorization of the Adults in a Small Town

	Employed	Unemployed	Total
Male	460	40	500
Female	140	260	400
Total	600	300	900

Conditional Probability (6/31)

Example2:

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being produced. These strips can be defective in two ways, length and nature of texture. It is known from historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests. If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

Conditional Probability (7/31)

Example2:

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being produced. These can be defective in two ways, length and nature of texture. It is known from historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests. If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

$P(L) = 0.1$

Conditional Probability (7/31)

Example2:

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being produced. These strips are defective in two ways, length and nature of texture. It is known from historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests. If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

$P(L) = 0.1$

$P(T) = 0.05$

Conditional Probability (7/31)

Example2:

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being produced. These strips are defective in two ways, length and nature of texture. It is known from historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests. If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

$P(L) = 0.1$

$P(T) = 0.05$

$P(L \cap T) = 0.008$

Conditional Probability (7/31)

Example2:

Consider an industrial process in the textile industry in which strips of a particular type of cloth are being pr[illegible]. These strip[illegible]fective in t[illegible]ngth and nature of [illegible]re. It is [illegible]rom historical information on the process that 10% of strips fail the length test, 5% fail the texture test, and only 0.8% fail both tests.

$P(L) = 0.1$

$P(T) = 0.05$

$P(L \cap T) = 0.008$

If a strip is selected randomly from the process and a quick measurement identifies it as failing the length test, what is the probability that it is texture defective?

$$P(T|L) = 0.008/0.1 = 0.08$$

Conditional Probability (8/31)

Example3:

A die is rolled twice. What is the probability that the sum equal 10, if you know that the 1st element equal 6?

Conditional Probability (9/31)

Example3:

A die is rolled twice. What is the probability that the sum equal 10, if you know that the 1st element equal 6?

Solution: $A = \{46, 55, 64\}$, $B = \{61, 62, 63, 64, 65, 66\}$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Conditional Probability (9/31)

Example3:

A die is rolled twice. What is the probability that the sum equal 10, if you know that the 1st element equal 6?

Solution: $A = \{46, 55, 64\}$, $B = \{61, 62, 63, 64, 65, 66\}$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$P(A) = 3/36$
$P(B) = 6/36$

$P(A \cap B) = 1/36$

$(A \cap B) = \{64\}$

Conditional Probability (9/31)

Example3:

A die is rolled twice. What is the probability that the sum equal 10, if you know that the 1st element equal 6?

Solution: $A = \{46, 55, 64\}$, $B = \{61, 62, 63, 64, 65, 66\}$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{1/36}{6/36} = 1/6$$

$P(A) = 3/36$
$P(B) = 6/36$

$P(A \cap B) = 1/36$

$(A \cap B) = \{64\}$

Conditional Probability (10/31)

Disjoint (or mutually exclusive):

S is a sample space, A, B are two events

$A, B \subseteq S$ and A, B **Disjoint** *or* **mutually exclusive**

$$\therefore P(A \cap B) = 0$$

$$\therefore P(A \cup B) = P(A) + P(B)$$

$$\therefore P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{P(B)} = 0$$

Conditional Probability (11/31)

Independence:

S is a sample space, A, B are two events
$A, B \subseteq S$ and A, B are **independent**

$$\therefore P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A) * P(B)}{P(B)} = P(A)$$

$$\therefore P(A \cap B) = P(A) * P(B)$$

$$\therefore P(A \cup B) = P(A) + P(B) - P(A) * P(B)$$

Conditional Probability (12/31)

Example4:

- If $P(A) = 0.2$, $P(B) = 0.3$ determine the following probabilities:

If A, and B are **disjoint** (mutually exclusive)

1) $P(A \cap B)$
2) $P(A \cup B)$
3) $P(A|B)$

Conditional Probability (13/31)

Example4:

- If $P(A) = 0.2$, $P(B) = 0.3$ determine the following probabilities:

If A, and B are **disjoint** (mutually exclusive)

Solution:

$$P(A \cap B) = 0$$

$$P(A \cup B) = P(A) + P(B) = 0.2 + 0.3 = 0.5$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0}{P(B)} = 0$$

Conditional Probability (14/31)

Example5:

- If $P(A) = 0.2$, $P(B) = 0.3$ determine the following probabilities:

If A, and B are **independent**

1) $P(A \cap B)$
2) $P(A \cup B)$
3) $P(A|B)$

Conditional Probability (15/31)

Example5:

- If $P(A) = 0.2$, $P(B) = 0.3$ determine the following probabilities:

If A, and B are **independent**

Solution:

$$P(A \cap B) = P(A) * P(B) = 0.2 * 03 = 0.06$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.5 - 0.06 = 0.44$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A) * P(B)}{P(B)} = P(A) = 0.2$$

Conditional Probability (16/31)

Example6:

The following circuit operates only if there is a path of functional devices from left to right. The probability that each device functions is shown on the graph. Assume that devices fail independently. What is the probability that the circuit operates?

Conditional Probability (17/31)

Example6:

The following circuit operates only if there is a path of functional devices from left to right. The probability that each device functions is shown on the graph. Assume that devices fail independently. What is the probability that the circuit operates?

Let L and R denote the events that the left and right devices operate, respectively.

$$P(L \cap R) = P(L)P(R) = 0.80(0.90) = 0.72$$

Conditional Probability (18/31)

Example7:

The following circuit operates only if there is a path of functional devices from left to right. The probability that each device functions is shown on the graph. Assume that devices fail independently. What is the probability that the circuit operates?



Conditional Probability (19/31)

Example7:

What is the probability that the circuit operates?



Let T and B denote the events that the top and bottom devices operate, respectively.

$$P(T \cup B) = P(T) + P(B) - P(T)P(B)$$

$$= 0.95 + 0.90 - (0.95)(0.90) = 0.995$$

Conditional Probability (20/31)

Example7:

Another Solution

What is the probability that the circuit operates?



$$P(T \cup B) = 1 - P(T \cup B)'$$

$$= 1 - P(T' \cap B') = 1 - P(T')P(B')$$

$$= 1 - (0.05)(0.10) = 0.995$$

Conditional Probability (21/31)

Example8:

The following circuit operates only if there is a path of functional devices from left to right. The probability that each device functions is shown on the graph. Assume that devices fail independently. What is the probability that the circuit operates?



Conditional Probability (22/31)

Example8:



$$= 1 - (0.10)(0.10)(0.10) = 0.999$$

Conditional Probability (23/31)

Example8:



$$= 1 - (0.05)(0.05) = 0.9975$$

Conditional Probability (24/31)

Example8:



$$= (0.999)(0.9975)(0.99) = 0.9865$$

Conditional Probability (25/31)

Multiplication Rule:

$$P(A|B) = \frac{P(A \cap B)}{P(B)} \quad , \qquad for \quad P(B) > 0$$

$$P(B|A) = \frac{P(A \cap B)}{P(A)} \quad , \qquad for \quad P(A) > 0$$

$$\boldsymbol{P(A \cap B) = P(A|B)P(B) = P(B|A)P(A)}$$

Conditional Probability (26/31)

Example9:

One bag contains 4 white balls and 3 black balls, and a second bag contains 3 white balls and 5 black balls. One ball is drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from the second bag is black?

Conditional Probability (27/31)

Example9:

One bag contains 4 white balls and 3 black balls, and a second bag contains 3 white balls and 5 black balls. One ball is drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from the second bag is black?





Solution:

B_1: Black from bag#1

W_1: White from bag#1

B_2: Black from bag#2

W_2: White from bag#2

Conditional Probability (27/31)

Example9: B_1 and B_2 or W_1 and B_2

One bag contains 4 white balls and 3 black balls, and a second bag contains 3 white balls and 5 black balls. One ball is drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from the second bag is black?





Solution:

B_1: Black from bag#1

W_1: White from bag#1

B_2: Black from bag#2

W_2: White from bag#2

Conditional Probability (28/31)

Example9: B_1 and B_2 or W_1 and B_2

$$P(B_1 \cap B_2) = P(B_2|B_1)P(B_1) = \left(\frac{6}{9}\right)\left(\frac{3}{7}\right)$$





Conditional Probability (29/31)

Example9: B_1 and B_2 or W_1 and B_2

$$P(W_1 \cap B_2) = P(B_2|W_1)P(W_1) = \left(\frac{5}{9}\right)\left(\frac{4}{7}\right)$$



Conditional Probability (30/31)

Example9: B_1 and B_2 or W_1 and B_2 **Disjoint**

$$P(B_1 \cap B_2) = P(B_2|B_1)P(B_1) = \left(\frac{6}{9}\right)\left(\frac{3}{7}\right)$$

$$P(W_1 \cap B_2) = P(B_2|W_1)P(W_1) = \left(\frac{5}{9}\right)\left(\frac{4}{7}\right)$$

What is the probability that a ball now drawn from the second bag is black = $\left(\frac{6}{9}\right)\left(\frac{3}{7}\right) + \left(\frac{5}{9}\right)\left(\frac{4}{7}\right) = \frac{38}{63}$

Conditional Probability (31/31)

Example9: B_1 and B_2 or W_1 and B_2 **Disjoint**



Video Lectures

All Lectures: https://www.youtube.com/playlist?list=PLxIvc-MGOs6gW9SgkmoxE5w9vQkID1_r-

Lecture #3: https://www.youtube.com/watch?v=raeVQxzY7iE&list=PLxIvc-MGOs6gW9SgkmoxE5w9vQkID1_r-&index=3

Thank You

Dr. Ahmed Hagag
ahagag@fci.bu.edu.eg